


# NATIONAL CHILDHOOD IMMUNISATION SCHEDULE



## Notes:

BCG	Bacillus Calmette-Guérin
HepB	Hepatitis B vaccine
DTaP	Paediatric diphtheria and tetanus toxoid and acellular pertussis vaccine
Tdap	Tetanus toxoid, reduced diphtheria toxoid and acellular pertussis vaccine
IPV	Inactivated polio vaccine
OPV	Oral polio vaccine
Hib	Haemophilus influenza type b vaccine
MMR	Measles, mumps and rubella vaccine
HPV	Human Papillomavirus which can cause cervical cancer and genital warts in women

PCV	Pneumococcal conjugate vaccine
D1/D2/D3	1 <sup>st</sup> dose, 2 <sup>nd</sup> dose, 3 <sup>rd</sup> dose
B1/B2	1 <sup>st</sup> booster, 2 <sup>nd</sup> booster
^	Primary 5
#	3 <sup>rd</sup> dose of HepB can be given between 5-6 months with the 3 <sup>rd</sup> dose of DTaP and IPV for the convenience of parents
##	2 <sup>nd</sup> dose of MMR can be given between 15-18 months
*	Recommended for females 9 to 26 years; three doses are required at intervals of 0, 2, 6 months