National Childhood Immunisation Schedule (NCIS) (From birth to age 17 years, effective from 1 November 2020)

Vaccine	Birth	2 months	4 months	6 months	12 months	15 months	18 months	2-4 years	5-9 years	10-11 years	12-13 years	13-14 years	15-17 years
Bacillus Calmette-Guérin (BCG)	D1												
Hepatitis B (HepB)	D1	D2		D3									
Diphtheria, tetanus and acellular pertussis (paediatric) (DTaP)		D1	D2	D3			B1						
Tetanus, reduced diphtheria and acellular pertussis (Tdap)										B2			
Inactivated poliovirus (IPV)		D1	D2	D3			B1			B2			
Haemophilus influenzae type b (Hib)		D1	D2	D3			B1						
Pneumococcal conjugate (PCV10 or PCV13)			D1	D2	B1								
Pneumococcal polysaccharide (PPSV23)									ne or two doses for children and adolescents age 2-17 years ith specific medical condition or indication.				
Measles, mumps and rubella (MMR)					D1	D2							
Varicella (VAR)					D1	D2							
Human papillomavirus (HPV2 or HPV4)											D1 (females)	D2 (females)	
Influenza (INF)				Annual vaccination or per season for <u>all children</u> age 6 months to <5 years (6-59 months).					Annual vaccination or per season for children and adolescents age 5-17 years with specific medical condition or indication.				



Recommended ages and doses for all children



Recommended for children with specific medical conditions or indications

Footnotes:

- D1, D2, D3: Dose 1, dose 2, dose 3
- B1, B2: Booster 1, booster 2
- 10-11, 12-13, 13-14 years: Primary 5, Secondary 1, Secondary 2 (Tdap, IPV, HPV (for females) and MMR (as catch-up) vaccines are provided as part of Health Promotion Board's school-based vaccination programme)
- HepB: Doses 2 and 3 are recommended to be given as part of the 6-in-1 vaccine at 2 and 6 months, respectively
- MMR: Only the dose 2 is recommended to be given as part of the MMRV vaccine

Speak to your doctor about your recommended vaccines. Visit vaccinesprotect.gov.sg for more information.